


Measuring altered movement patterns of musicians - Accuracy of a Motion Capturing System outside laboratory conditions

Nikolaus Ballenberger, Dirk Möller, Eduard Wolf, Christoff Zalpour



Background

- Use of **camera-based movement analysis** systems in medicine, kinesiology, biomechanics and in music health sciences
 - Identification of **altered/disturbed movement** patterns which lead to an **increased risk** of injury/illness for musicians (Davis,1997; Dixon,2011)
- Research project: “**Musikphysioanalysis**”
 - Do musicians with playing-related musculoskeletal complaints show altered movement patterns?

Background

- **high accuracy** of measurements under lab conditions (Everaert,1998; Windolf,2008)
 - **Advantage:** Motion capturing systems are **not restricted** to lab use
 - Measurement **under mobile conditions** in familiar/natural surroundings e.g. when rehearsing....
- However, no data available of measurement accuracy out of lab



HOCHSCHULE
OSNABRÜCK

UNIVERSITY OF APPLIED SCIENCES

Aim of study

To investigate whether **mobile use** of a marker based motion capturing system **impairs** measurement **accuracy** by means of a **two “pragmatic” experiments** (accuracy = absence of systematic and random error)



HOCHSCHULE
OSNABRÜCK

UNIVERSITY OF APPLIED SCIENCES

Methods: Experiment 1

Wand swung in measurement volume (60 sec) in **three scenarios:**

- Out of lab (8m x 4m x 2.5m)
- Out of lab (5m x 3m x 2.5m)
- Lab (10m x 8m x 2.5m)

→ Compare measured distances to **true value (600.8 mm)**



Methods: Experiment 2


musician playing **chromatic scale** with „**marked**“ violin in in **three scenarios**:

- Out of lab (8m x 4m x 2.5m)
- Out of lab (5m x 3m x 2.5m)
- Lab (10m x 8m x 2.5m)

Comparison to „**true**“ values → under static conditions:

→ distances (true values)

→ Angles (true values)



Methods: Calculating measurement error

- Calculation all **possibles distances/angles** between all markers
- Calculation of **deviation** from **true value** (measurement error) for each of the 9000 measurements for **each marker** separately
- processed by **moving avarage**



HOCHSCHULE
OSNABRÜCK

UNIVERSITY OF APPLIED SCIENCES

Methods: Calculating measurement error

k denotes the number of markers m and n the number of measurements i

Deviation from true value of one single measurement = $x_{mi} - \text{true value}_m$

$$\text{Total error of marker}_m \text{ TE}_m = \sqrt{\frac{1}{n} \sum_{i=1}^n (x_{mi} - \text{true value}_m)^2}$$

→ Decomposition of TE_m into **systematic** and **random** error for each marker

$$\text{Random error of marker}_m \text{ RE}_m = \sqrt{\frac{1}{n} \sum_{i=1}^n (x_{mi} - \bar{x}_m)^2}$$

$$\text{Systematic error of marker}_m \text{ SE}_m = \bar{x}_m - \text{true value}_m$$

Overall error across all markers → decomposition into: **squared average random error** of all markers and **variance of systematic errors** of all markers and **average systematic error** of all markers

$$\sqrt{\overline{\text{RE}}^2 + \frac{1}{k} \sum_{m=1}^k (\text{SE}_m - \overline{\text{SE}})^2 + \overline{\text{SE}}^2} \quad \text{or} \quad \sqrt{\frac{1}{kn} \sum_{m=1}^k \sum_{i=1}^n (x_{mi} - \text{true value}_m)^2}$$




HOCHSCHULE
OSNABRÜCK

UNIVERSITY OF APPLIED SCIENCES

Methods: Calculating measurement error (Graphical illustration)

Distribution of measurement error during a 60 sec record)

Random error = dispersion
around measured mean




systematic error = mean deviation from zero
(no deviation from true value)

Methods: Calculating measurement error (Graphical illustration)


Total error: pooling errors across **all** markers

Random error = dispersion
around measured mean




systematic error = mean deviation from zero
(no deviation from true value)


Deviation (distance) from true value depending (experiment 1 with wand)



Deviation (distance) from true value depending (experiment 1 with wand)



Deviation (distance) from true value depending on scenario and marker (experiment 2 with violin)




Out of lab with large volume Out of lab with small volume

Lab

Deviation (distance) from true value depending on scenario and marker (experiment 2 with violin)

No reduction in accuracy so far between lab and out of lab....

frequency




Out of lab with large volume


Out of lab with small volume

Lab

Deviation (distance) from true value depending on scenario and marker (experiment 2 with violin)




Deviation (distance) from true value depending on scenario and marker (experiment 2 with violin)



Out of lab with large volume Out of lab with small volume

Lab

Deviation (distance) from true value depending on scenario and marker (experiment 2 with violin)




Deviation (distance) from true value depending on scenario and marker (experiment 2 with violin)

\overline{RE}	\overline{SE}	SD (SE)
0.25	-0.23	0.34
Overall TE		
0.49		

\overline{RE}	\overline{SE}	SD (SE)
0.21	-0.21	0.26
Overall TE		
0.41		

\overline{RE}	\overline{SE}	SD (SE)
0.21	-0.50	4.16
Overall TE		
4.19		



frequency

-5

0

deviation from true value (mm)

5

0

deviation from true value (mm)

-5

0


deviation from true value (mm)

5


Out of lab with large volume Out of lab with small volume

Lab


Deviation (distance) from true value depending on scenario and marker (experiment 2 with violin)





Deviation (distance) from true value depending on scenario and marker (experiment 2 with violin)





What is the problem?





What is the problem?




Merging of markers




no reduction of accuracy under mobile conditions
but due to measurement volume




Deviation (distance) from true value depending on scenario and marker (experiment 2 with violin)




Out of lab (8m x 4m x 2.5m)



Out of lab (5m x 3m x 2.5m)



Lab (10m x 8m x 2.5m)



Discussion/Conclusion

- no loss of accuracy out of lab
- **Large** volume (long distance of cameras) leads to **merging** of close markers
- **No conclusion** about accuracy when markers are **changing positions**
→ **however**: no assessment of accuracy possible due to **missing gold standard**
- **no guarantee** of accuracy with calibration by **wand**



HOCHSCHULE
OSNABRÜCK

UNIVERSITY OF APPLIED SCIENCES

Discussion/Conclusion

Message :

- Measureing **out of lab** is without reducing **accuracy**
- **BUT:** be aware of **optimal volume** espacially with **close markers!**



HOCHSCHULE
OSNABRÜCK

UNIVERSITY OF APPLIED SCIENCES

References

- Mitchell HL. Applications of digital photogrammetry to medical investigations. ISPRS Journal of Photogrammetry and Remote Sensing 1995;50(3):27. doi:10.1016/0924-2716(95)91288-U.
- Davis RB. Reflections on clinical gait analysis. J Electromyogr Kinesiol 1997;7(4):251. ENG.
- Dixon S. Principles of biomechanics and their use in the analysis of injuries and technique. In: Hutson MA, Speed C, editors. Sports injuries. Oxford u.a.: Oxford Univ. Press; 2011. p. 96–102. ISBN: 9780199533909.
- Everaert DG, Spaepen AJ, Wouters MJ, Stappaerts KH, Oostendorp RA. Measuring small linear displacements with a three-dimensional video motion analysis system: determining its accuracy and precision. Arch Phys Med Rehabil 1999;80(9):1082. eng.
- Richards JG. The measurement of human motion, A comparison of commercially available systems. Human Movement Science 1999;18(5):589. doi:10.1016/S0167-9457(99)00023-8.
- Maletsky LP, Sun J, Morton NA. Accuracy of an optical active-marker system to track the relative motion of rigid bodies. J Biomech 2007;40(3):682. eng. doi:10.1016/j.jbiomech.2006.01.017.
- Windolf M, Gotzen N, Morlock M. Systematic accuracy and precision analysis of video motion capturing systems--exemplified on the Vicon-460 system. J Biomech 2008;41(12):2776. eng. doi:10.1016/j.jbiomech.2008.06.024.



HOCHSCHULE
OSNABRÜCK

UNIVERSITY OF APPLIED SCIENCES

MUSICPHYSIO 2018

2nd World Congress on
MUSICIAN'S
PHYSIOTHERAPY

SEP 06 - 08 // 18
UNIVERSITY OF APPLIED SCIENCES
OSNABRUECK, GERMANY



Lectures

Prof. Dr. Ulrik Roijezon (*Lulea University, Sweden*)
Posture, Motor Control & Performance

Robert Kuizenga (*Counter Tenor, PT, Amsterdam*)
Physiotherapy & Voice

Prof. Dr. Bronwen J. Ackermann (*University of Sydney, Australia*)
Musicians' Physiotherapy: Past, Present and Future

Prof. Dr. med. Eckart Altenmüller (*hmtm Hannover, Germany*)
***Movement Disorders in Musicians:
Differential Diagnosis and Treatment Options***



HOCHSCHULE OSNABRÜCK
UNIVERSITY OF APPLIED SCIENCES

Workshops

Ed Blake (*Physio ED, London*)
Orofacial and Neck Assessment & Management

Dr. Sonia Ranelli (*Curtin University, Perth*)
***Hand and Upper Limb Problems in Musicians:
Physical and Sensory Assessment in Musicians
and Implications for Management***

Dr. Jan Dommerholt (*Bethesda, Baltimore*)
Management of Myofascial Pain

...and many more